

LESKO® DIMENSION* 0.10% Plus Fertilizer

NOT FOR USE on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

In New York State this product may only be used by commercial applicators and at no more than 500 lb (0.5 lb of active ingredient) per acre per year (or 11.5 lb product per 1,000 ft² sq per year) and is prohibited from use in Nas-sau and Suffolk Counties.

ACTIVE INGREDIENT: LESKO® Poly Plus® Polymer Coated Urea to provide a uniform growth with extended nitrogen feeding.

NET WEIGHT: 50 lb (22.7 kg) net weight

DIMENSIONS: 35" x 16" x 16" (914 mm x 406 mm x 406 mm)

NET CONTENTS: 99.90% LESKO® Polymer Coated Urea, 0.10% Poly Plus® Polymer Coated Urea

PRECAUTIONS: Before using this product, read the Use Precautions, Warranty Statements, Directions for Use, and the Storage and Disposal Instructions. If the Warranty statements are not acceptable, return the product unused, and the

within thirty days of purchase to the place of purchase.

Read the entire label before using this product. Use only according to label instructions.

Product protected by U.S. Patent No. 4,892,184. Other patents pending.

KEEP OUT OF REACH OF CHILDREN

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

If on skin or clothing: Take off contaminated clothing. Raise skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! CAUSES EYE IRRITATION, HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wear protective eyewear such as goggles, face shield or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Treated areas do not contain water when weather conditions favor drift from

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass, other annual grasses, and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed good root system and a uniform stand, and have received at least two mowings prior to lawn and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury. This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrasses species that have been determined to be tolerant to applications of this product. When applied as directed under these use directions, the following established turfgrasses are tolerant to this product:

Common Name	Scientific Name
Bentgrass, creeping	<i>Agrostis palustris</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, tall	<i>Festuca arundinacea</i>
Fescue, fine	<i>Festuca rubra</i>
Fescue, hard	<i>Festuca arundinacea</i>
Kygrass, perennial	<i>Lolium perenne</i>

Warm-Season Grasses

Cool-Season Grasses

DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis* tenuis) varieties:

Common Name	Scientific Name
Bahagrass	<i>Paspalum notatum</i>
Bermudagrass	<i>Cynodon dactylon</i>
Buffalograss**	<i>Bouteloua dactyloides</i>
Caregrass	<i>Axonopus affinis</i>
Centipedegrass	<i>Eranthis ciliaris</i>
Kikuyugrass	<i>Pennisetum clandestinum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>

**DO NOT apply this product on seedling Buffalograss in the spring of the first year of establishment until the turf-dmF-93; "Highlight", "Valo", "Jamestown", "Koker", "Majesta", "Mary", "Pannawm", "Tamera", "Tajana", "Waldorf", and "Wal-dorf". Not all varieties of Fine Fescue have been tested.

Use of this product on certain varieties of Creeping Bentgrass, such as "Cohasset", "Carmen", "Seaside", and "Washington" may result in undesirable turfgrass injury. Not all varieties of Creeping Bentgrass have been tested.

Fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bijart", "C-F-2", "Enjoy", "HF-93", "Highlight", "Valo", "Jamestown", "Koker", "Majesta", "Mary", "Pannawm", "Tamera", "Tajana", "Waldorf", and "Wal-dorf". Not all varieties of Fine Fescue have been tested.

For single applications made in the spring or early summer, this product should be applied at the appropriate rate according to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be pre-emergence or early post-emergence at the time of application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, turfgrass competitiveness, and the user's location within a region.

Use Program #1 for pre-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of pre-emergence crabgrass control. This program should not be used for early post-emergence crabgrass control.

Use Program #2 for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turf-grass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program #3 for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf-grass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or post-emergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1: Recommended Single Application Use Rates*

USE REGION	#1	#2	#3
North	2.86 lb/1,000 sq ft	5.73 lb/1,000 sq ft	5.73 lb/1,000 sq ft
Transition	4.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	8.49 lb/1,000 sq ft
South	5.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	11.47 lb/1,000 sq ft
Coastal South	5.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	11.47 lb/1,000 sq ft

Note: 5.73 lb per 1,000 sq ft is equal to 0.25 lb active ingredient per acre. Particle distribution at this rate may not provide adequate control or suppression. ‡ DO NOT apply more than 11.47 lb of this product per 1,000 sq ft per application, and no more than 34.41 lb of this product per year. Post-emergence control is limited: see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below. See map of the United States. All areas not designated below.

North: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NM, NV, OK, SC, TN, & TX. Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

** May require split or sequential applications for full control.

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after application. When treated lawn or ornamental turf areas are wetted or receive significant rainfall within a few days after application, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Other Grasses and Broadleaf Weeds

Application DIRECTIONS - Control of Crabgrass

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

NET WEIGHT: 50 lb (22.7 kg) net weight

DIMENSIONS: 35" x 16" x 16" (914 mm x 406 mm x 406 mm)

NET CONTENTS: 99.90% LESKO® Polymer Coated Urea, 0.10% Poly Plus® Polymer Coated Urea

PRECAUTIONS: Before using this product, read the Use Precautions, Warranty Statements, Directions for Use, and the Storage and Disposal Instructions. If the Warranty statements are not acceptable, return the product unused, and the

within thirty days of purchase to the place of purchase.

Read the entire label before using this product. Use only according to label instructions.

Product protected by U.S. Patent No. 4,892,184. Other patents pending.

KEEP OUT OF REACH OF CHILDREN

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.

If on skin or clothing: Take off contaminated clothing. Raise skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION! CAUSES EYE IRRITATION, HARMFUL IF INHALED. Avoid contact with eyes or clothing. Avoid breathing dust. Wear protective eyewear such as goggles, face shield or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Prolonged or frequently repeated skin contact while handling this material may cause allergic reaction in some individuals.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff from treated turf may adversely affect aquatic organisms in adjacent aquatic sites. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Treated areas do not contain water when weather conditions favor drift from

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

GENERAL INFORMATION

This product is a herbicide that provides control of crabgrass, other annual grasses, and broadleaf weeds in established lawns and ornamental turfs, including golf course fairways, roughs, and tee boxes. This product will not harm nearby established ornamentals when used according to label directions.

This product may be used on seeded, sodded, or sprigged lawns and ornamental turfs that are well-established. The grass must have developed good root system and a uniform stand, and have received at least two mowings prior to lawn and ornamental turfs that are not well-established, or on those that have been weakened by weather-, pest-, disease-, chemical-, or mechanical-related stress, may increase the chances for turf injury. This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrasses species that have been determined to be tolerant to applications of this product. When applied as directed under these use directions, the following established turfgrasses are tolerant to this product:

Common Name	Scientific Name
Bentgrass, creeping	<i>Agrostis palustris</i>
Bluegrass, Kentucky	<i>Poa pratensis</i>
Fescue, tall	<i>Festuca arundinacea</i>
Fescue, fine	<i>Festuca rubra</i>
Fescue, hard	<i>Festuca arundinacea</i>
Kygrass, perennial	<i>Lolium perenne</i>

Warm-Season Grasses

Cool-Season Grasses

DO NOT apply this product to Colonial Bentgrass (*Agrostis tenuis* tenuis) varieties:

Common Name	Scientific Name
Bahagrass	<i>Paspalum notatum</i>
Bermudagrass	<i>Cynodon dactylon</i>
Buffalograss**	<i>Bouteloua dactyloides</i>
Caregrass	<i>Axonopus affinis</i>
Centipedegrass	<i>Eranthis ciliaris</i>
Kikuyugrass	<i>Pennisetum clandestinum</i>
St. Augustinegrass	<i>Stenotaphrum secundatum</i>
Zoysiagrass	<i>Zoysia japonica</i>

**DO NOT apply this product on seedling Buffalograss in the spring of the first year of establishment until the turf-dmF-93; "Highlight", "Valo", "Jamestown", "Koker", "Majesta", "Mary", "Pannawm", "Tamera", "Tajana", "Waldorf", and "Wal-dorf". Not all varieties of Fine Fescue have been tested.

Use of this product on certain varieties of Creeping Bentgrass, such as "Cohasset", "Carmen", "Seaside", and "Washington" may result in undesirable turfgrass injury. Not all varieties of Creeping Bentgrass have been tested.

Fescue varieties have been found to be sensitive to this product: "Atlanta", "Banner", "Beauty", "Bijart", "C-F-2", "Enjoy", "HF-93", "Highlight", "Valo", "Jamestown", "Koker", "Majesta", "Mary", "Pannawm", "Tamera", "Tajana", "Waldorf", and "Wal-dorf". Not all varieties of Fine Fescue have been tested.

For single applications made in the spring or early summer, this product should be applied at the appropriate rate according to one of the three control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be pre-emergence or early post-emergence at the time of application. The duration of residual weed control provided by this product is directly related to the total rate applied, but will vary somewhat depending on weather, turfgrass competitiveness, and the user's location within a region.

Use Program #1 for pre-emergence control at sites where the turfgrass is cut high (e.g., homeowner lawns). This program provides 3-5 months of pre-emergence crabgrass control. This program should not be used for early post-emergence crabgrass control.

Use Program #2 for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways), and (b) turf-grass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut high (e.g., homeowner lawns).

Use Program #3 for pre-emergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and (b) turf-grass maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of pre-emergence crabgrass control. This program may also be used for early post-emergence control at sites where turfgrass is cut low (e.g., golf fairways).

Subsequent, sequential pre- and/or post-emergence applications should be made where longer periods of control are desired.

Where split fertilizer applications are recommended, the rates in Table 1 may be split across two applications made 6-10 weeks apart and prior to crabgrass emergence.

Table 1: Recommended Single Application Use Rates*

USE REGION	#1	#2	#3
North	2.86 lb/1,000 sq ft	5.73 lb/1,000 sq ft	5.73 lb/1,000 sq ft
Transition	4.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	8.49 lb/1,000 sq ft
South	5.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	11.47 lb/1,000 sq ft
Coastal South	5.73 lb/1,000 sq ft	8.49 lb/1,000 sq ft	11.47 lb/1,000 sq ft

Note: 5.73 lb per 1,000 sq ft is equal to 0.25 lb active ingredient per acre. Particle distribution at this rate may not provide adequate control or suppression. ‡ DO NOT apply more than 11.47 lb of this product per 1,000 sq ft per application, and no more than 34.41 lb of this product per year. Post-emergence control is limited: see "Crabgrass Control" and "Precautions" sections above. Regions include areas listed below. See map of the United States. All areas not designated below.

North: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, IN, OH, & coastal areas of CT, NY, & RI.

South: AL, AR, AZ, CA, GA, LA, MS, NM, NV, OK, SC, TN, & TX. Coastal South: HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

** May require split or sequential applications for full control.

This product can also be applied in the late summer or early fall (late August through November) at the "Program 3" use rates listed in Table 1 to provide control of crabgrass through the early part of the next spring. The fall application should be followed by an appropriate spring application to provide season-long control.

Tips for Improved Control

For best results, apply this product within a few days after mowing and delay mowing again for a few days after application. When treated lawn or ornamental turf areas are wetted or receive significant rainfall within a few days after application, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Other Grasses and Broadleaf Weeds

Application DIRECTIONS - Control of Crabgrass

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

Fall Applications

Used as directed for late summer or early fall use, for crabgrass control through the early part of the next spring, this product will also control (at the "Program 3" rates) the following weeds when applied prior to their emergence:

Also controlled at the "Program 2" rates:

NET WEIGHT: 50 lb (22.7 kg) net weight

DIMENSIONS: 35" x 16" x 16" (914 mm x 406 mm x 406 mm)

NET CONTENTS: 99.90% LESKO® Polymer Coated Urea, 0.10% Poly Plus® Polymer Coated Urea

PRECAUTIONS: Before using this product, read the Use Precautions, Warranty Statements, Directions for Use, and the Storage and Disposal Instructions. If the Warranty statements are not acceptable, return the product unused, and the

within thirty days of purchase to the place of purchase.

Read the entire label before using this product. Use only according